



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Fertility preservation and preimplantation genetic assessment for women with breast cancer

Citation for published version:

Sciorio, R & Anderson, RA 2019, 'Fertility preservation and preimplantation genetic assessment for women with breast cancer', *Cryobiology: International Journal of Low Temperature Biology and Medicine*.
<https://doi.org/10.1016/j.cryobiol.2019.12.001>

Digital Object Identifier (DOI):

[10.1016/j.cryobiol.2019.12.001](https://doi.org/10.1016/j.cryobiol.2019.12.001)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Cryobiology: International Journal of Low Temperature Biology and Medicine

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy


The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Journal Pre-proof

Fertility preservation and preimplantation genetic assessment for women with breast cancer

Romualdo Sciorio, Richard A. Anderson



PII: S0011-2240(19)30535-8

DOI: <https://doi.org/10.1016/j.cryobiol.2019.12.001>

Reference: YCRYO 4156

To appear in: *Cryobiology*

Received Date: 2 November 2019

Revised Date: 17 December 2019

Accepted Date: 17 December 2019

Please cite this article as: R. Sciorio, R.A. Anderson, Fertility preservation and preimplantation genetic assessment for women with breast cancer, *Cryobiology* (2020), doi: <https://doi.org/10.1016/j.cryobiol.2019.12.001>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier Inc.

Fertility preservation and preimplantation genetic assessment for women with breast cancer

Romualdo Sciorio^{1*}, Richard A. Anderson²

1. Edinburgh Assisted Conception Programme, EFREC, Royal Infirmary of Edinburgh, 51 Little France Crescent, Edinburgh, UK.

2. MRC Centre for Reproductive Health, The Queen's Medical Research Institute, The University of Edinburgh, 47 Little France Crescent, Edinburgh, UK.

***Corresponding author. E-mail address: sciorioromualdo@hotmail.com (R. Sciorio).**

Abstract

Breast cancer is the most common cancer diagnosed among reproductive aged women, and its treatment can compromise future fertility. Options for fertility preservation include oocyte or embryo cryopreservation after ovarian stimulation (OS), which are the most established choices and are applicable for adult women with cancer. Ovarian tissue freezing may also be appropriate, as it offers potentially the least delay. The recognition of the role of BRCA1 and BRCA2 mutations in some women has led to the involvement of preimplantation genetic diagnosis (PGD), recently renamed preimplantation genetic testing for monogenic disorder (PGT-M), whereby embryos are created by IVF and cell(s) are removed and genetically analyzed for specific disease-related mutations. PGT-M offers a valid option for women wishing to avoid transmission of the predisposition for hereditary breast cancer to their offspring. The aim of this paper is to provide an overview of the factors that influence fertility preservation in newly diagnosed breast cancer patients, and to illustrate the option of PGT-M to enable conception of an unaffected child.

KEYWORDS: Breast cancer, Ovarian stimulation, PGT-M, Fertility preservation, Oocyte cryopreservation, Embryo cryopreservation, Ovarian tissue cryopreservation.

Background

Breast cancer (BC) is the most common cancer in women of reproductive age, with more than 10% of new cases diagnosed in women younger than the age of 40 years [41]. Currently, with the social trend to delaying motherhood until later in life, there are an increasing number of women who have

not completed childbearing at the time of cancer diagnosis, and therefore are likely to desire pregnancy following the chemotherapy [56]. In 2018, has been calculated that 2.1 million new cases of BC were diagnosed worldwide [12]. For many years, BC has been considered the most important cancer in reproductively-aged women, both in terms of incidence and mortality. However, for a range of reasons including improved screening methods and therapies, the number of deaths has been decreasing. Whereas in 2009, estimated deaths were 21.1% of estimated new cases, they were 15.4% in 2018, with a reduction of 27% over the last decade [22]. However, a potential side effect is the loss of fertility or impaired reproductive function [81]. Additionally, women with hormone receptor positive disease may also be advised to take hormonal therapy for up to 10 years after chemotherapy. This, also impact on the complexity of reproductive choices they have to make, facing declining fertility through increasing age as well as from effects of chemotherapy. Fertility concerns among young cancer patients have an important role in determining quality of life [69]. At the time diagnosis, about half of young women are concerned about becoming infertile or having reduced reproductive function after BC treatment, and while a survey 5 years ago indicated that only a small minority of 10% take up fertility preservation (FP) options [80], this proportion is increasing. There have also been concerns about whether a subsequent pregnancy may increase the chance of recurrence of breast cancer, but it is now clear that this is not the case [44].

Fertility Preservation: Available Options

The many advances in assisted reproductive technologies (ART) over the forty years since its introduction include the development of methods and strategies for FP in women with BC and other conditions whose treatment risks their future fertility, before initiation of anti-cancer therapy. These include cryopreservation of oocytes or embryos after OS, or ovarian tissue cryopreservation (OTC) [3, 27]. The recognition of the role of mutations in the BRCA1 and BRCA2 genes in the aetiology of breast and other cancers in some women also introduces consideration of the use of PGT-M in women with these genetic mutations in order to avoid transmitting the mutation to their offspring. These considerations have raised some concerns, but the possible health and psychological consequences of this particular condition are considered to justify its use [86]. These complex issues will be occurring at a time of great stress and uncertainty to patients in the immediate

aftermath of a new diagnosis. This and the very limited time available for discussion, decisions and potential interventions requires excellent lines of communication between the oncology setting and reproductive medicine. This review discusses the available methods for FP in women with breast cancer, and the role of PGT-M in this context. Protection of the ovary from chemotherapy-induced damage has also been the subject of significant investigation. This has recently been reviewed by Spears and colleagues [87], but of particular importance to women with breast cancer is the demonstration, now confirmed in several large RCTs, that administration of GnRH agonists during chemotherapy for breast cancer reduces the risk of premature ovarian insufficiency (POI). This has been subject to recent meta-analysis [45], thus will not be discussed in detail here. However it is important to recognize that while there seems good evidence regarding risk of POI, whether there is an increased chance of a subsequent pregnancy is unclear, and this approach should not be regarded as an effective form of FP where other interventions are possible.

Risks to fertility in breast cancer patients

Advances in chemotherapy and anti-cancer treatment have resulted in higher survival rates among cancer patients. The most common malignancy in adult women is breast cancer, affecting one in nine women [88]: the five-year survival rate for women treated for breast cancer in the UK is more than 80% [63]. Unfortunately, a side effect of chemotherapeutic drugs is the risk of developing POI, which is dependent on various factors. Most important is the chemotherapy regimen used and the drug doses: the alkylating agents are particularly gonadotoxic, but taxanes also have a negative effect [51]. The age of the patient is also important, as older women have a much higher reported incidence of POI after treatment, compared to the younger women [70, 49]. It is also clear that pre-chemotherapy ovarian reserve, as reflecting in serum concentrations of anti-Mullerian hormone (AMH) are also predictive of long-term ovarian function. This has been demonstrated in several prospective studies in women with BC [6, 7] also showing the interaction with age [89]. Pretreatment antral follicle count (AFC) may also be predictive, but there are few data clarifying this [90].

91

92 Effect of chemotherapy

93 Chemotherapy can have two different effects on ovarian function. The first is immediate, during or
94 following the treatment, with loss of the growing follicle population resulting in amenorrhea.
95 However, if sufficient primordial follicles remain in the resting pool upon the cessation of
96 treatment, the population of growing follicles will then be restored, and menses resume. In contrast,
97 the second is a longer term effect, caused by the depletion of the primordial follicle pool, and results
98 in a shortened reproductive lifespan and POI. If there is only partial loss of primordial follicles, this
99 longer term effect may not manifest itself until years following treatment [58]. Where the reduction
100 in the primordial follicle pool is near complete, the effect is acute, and the patient undergoes
101 immediate POI [70]. This results from the primordial pool of follicles being formed before birth,
102 such that at birth, the ovary has a fixed amount of oocytes. Primordial follicles are continuously
103 recruited out of the resting pool and activated to grow, but from each cohort of this follicles, only
104 very few will go through to the pre-ovulatory stage and eventually only one will ovulate: the
105 majority of follicles become atretic and will die at some point during development [36].
106 Chemotherapeutic agents can directly affect the resting pool of primordial follicles or the growing
107 follicles. The loss of the growing population of follicles may lead to increased activation of
108 primordial follicles and so the accelerated loss of that reserve. Chemotherapeutic agents target not
109 only the germ cells, but also the somatic cells. Granulosa cells surround the oocyte proliferate
110 during follicle maturation. Given the essential nature of the contact and communication between the
111 oocyte and the granulosa cells, damage to granulosa cells will result in indirect damage to the
112 oocyte, leading to follicle loss (Figure 1) [58].

113 It is difficult to predict the exact risk for future fertility. A population-based analysis of pregnancy
114 after cancer showed that women with breast cancer diagnosis before the age of 40 had a markedly
115 reduced chance of post-cancer pregnancy compared to age-matched controls, with a standardized
116 incidence ratio of 0.39 (95% confidence interval 0.36-0.42), but also that there have been significant
117 improvements in the chances of a post-cancer pregnancy over recent years (Figure 2) [4]. As stated
118 earlier, the gonadotoxic effect of chemotherapy is directly associated to female age at the time of
119 treatment and depends considerably on the agent used and the duration of treatment [51, 55]. With

reference to agents commonly used for breast cancer, alkylating agents have the strongest gonadotoxic potential. These agents, directly affect cell proliferation and primordial follicles [9], and promote cell apoptosis and follicle depletion [55]. Cyclophosphamide is one of the most effective drugs used for BC, is also the one of the most investigated compound in connection with gonadal toxicity: the risk of amenorrhea is high, and there is a four-fold higher risk of developing POI as compared with other agents [48, 51]. A high risk of amenorrhea, particularly in women in their later reproductive years, is also associated with other drugs such as fluorouracil, epirubicin and fluorouraciloxorubicin, which are often used in women with breast cancer. Taxanes cause an intermediate ovarian damage, whereas methotrexate and 5-fluorouracil are associated with a lower toxicity risk [25, 48]. Limited clinical data are currently available regarding newer agents such as trastuzumab, bevacizumab, and cetuximab [91]. Abusief and colleagues [29] suggested that trastuzumab might not induce amenorrhea in premenopausal women with breast cancer. However, further studies are needed to clarify the effect of these agents on ovarian function.

Oocyte Cryopreservation: from slow-freezing protocol

In the last decades, the cryopreservation of mature oocyte has become an established procedure in ART, and represents a safe and effective method for patients wishing to preserve their fertility [64, 97]. Oocyte quality is one of the most important factor influencing the vitrification-warming survival rate, and the subsequent fertilization and embryo development [31]. Cryopreservation involves freezing cells and subsequent storage in liquid nitrogen or its vapour at -196 °C. The first birth from a cryopreserved oocyte was reported in Australia in 1986, using a slow-freezing procedure [16]. Oocytes are extremely difficult cells to freeze successfully, mainly due to the large size cell, and the high content of water which during the freezing process might be converted to intracellular ice, which can induce damage and cell death [68]. Early studies highlighted difficulties in predicting the membrane permeability characteristics of human oocytes along with other biophysical components [29]. Several studies reported the negative effects of cryopreservation on the stability of microtubules and the spindle in mammalian oocytes [72]. In addition, zona pellucida (ZP) hardening after cryopreservation was reported as an extra complication from the freezing

process [97] although this can be overcome by the use of intracytoplasmic sperm injection (ICSI) [73]. Other possible injuries resulting from cooling and warming procedures include DNA fragmentation [33], damage to intracellular organelles [38] and epigenetic risks [99].

To Vitrification

A massive breakthrough in ART cryopreservation was reported with the introduction of “vitrification” in the late 1990s [43]. Vitrification was proposed as an alternative to the slow-freezing technique for human oocytes and embryos and was expected to give superior success rates in terms of cryo-survival and pregnancy outcomes. The Human Fertilisation and Embryology Authority (HFEA) has allowed the use of frozen oocytes for infertility treatment in the UK since 2000 [98] and the American Society for Reproductive Medicine (ASRM) in 2013 removed the experimental label applied to oocyte freezing [74] following randomized controlled studies [18, 77] which reported that IVF using vitrified-warmed oocytes could produce similar pregnancy outcomes to IVF with fresh oocytes. A recent meta-analysis confirmed that results from vitrification are superior to those achieved with slow freezing protocols [78]. An important consideration to make is the choice of the carrier used for vitrification, especially in terms of whether liquid nitrogen comes in contact with the droplet containing the embryo (open vitrification) or not (closed vitrification). The issue with open vitrification is that liquid nitrogen itself can contain microbes or pathogens, therefore concerns have been raised over the sterility of open systems due to potential cross contamination between the vitrification sample and liquid nitrogen [10]. Published studies have shown that closed vitrification devices can be used for successful cryopreservation of human embryos [82, 83, 96]. While some IVF scientists remain concerned that closed systems may reduce the survival rates, in the UK 75% of clinics use closed rather than open devices for vitrification [13].

Oocyte cryopreservation in cancer patients

The developments in oocyte cryopreservation described above can be considered a major advance in FP. Prior to the development of vitrification, slow freezing of oocytes had a very low success rate, and the more effective option of embryo cryopreservation was only available to women with a

178 partner, other than with the use of donated sperm. Cryopreservation of immature oocytes with
179 subsequent in vitro maturation is a potential option but still considered experimental [47], thus in
180 this section cryopreservation of mature oocytes (ie at metaphase II, MII) only will be discussed. A
181 key aspect of this approach is the need for OS, which takes at least 2 weeks, despite the
182 development of 'random start' protocols to minimise delay. These involve the administration of
183 FSH to stimulate multi-follicular development, which can be started at any stage in the menstrual
184 cycle, with co-administration of GnRH antagonists to prevent premature ovulation [23]. In general,
185 women with breast cancer respond to OS with the number of mature oocytes collected that would
186 be expected based on their age and pretreatment ovarian assessment [75]. Exposure to
187 supraphysiological levels of estrogen as a result of OS, albeit briefly, may be a particular risk for
188 patients with a hormone receptive cancer, and the aromatase inhibitor letrozole is widely used to
189 minimise this [94] without apparent detrimental effect on the ovarian response or the quality of the
190 oocytes recovered. Oktay and co-workers [64] analyzed the efficacy of oocyte cryopreservation by
191 vitrification in a meta-analysis, and reported live birth rates per oocyte warmed of 6.6%. A recent
192 study investigated the pregnancy outcome in fertility preservation after oocyte freezing for age-
193 related fertility decline and for patients before cancer treatment. This showed that overall oocyte
194 survival was comparable between the two groups, but implantation, ongoing pregnancy and live
195 birth rates were lower in cancer patients [20]. A live birth rate of 61.9% was reported from 12
196 cryopreserved oocytes in women ≤ 35 years and of 43.4% from 10 oocytes in those > 35 years, illustrating the
197 importance of both the number of oocytes that can be collected and cryopreserved (which of course declines
198 with age), and the decline in oocyte quality with age. Another aspect to be mentioned is the ideal
199 number of oocytes to freeze in order to obtain a pregnancy after warming. This is a critical point
200 that could be very useful and help clinicians to inform correctly their patients and plan their
201 treatments accordingly [34]. This aspect was investigated in a recent multicenter retrospective
202 study, included a total on 6,362 women who underwent to oocyte vitrification for FP, due to age-
203 related fertility decline (5,289 women) or for oncological reasons (1,172 women). The authors
204 reported an increased cumulative live birth rate from 15.8% with 5 oocytes to 32.0% with 8
205 oocytes. For younger patients (≤ 35 years old) 10 or 15 oocytes provided success rates of 42.8% and
206 69.8%. The highest cumulative live birth rate of 94.4% was obtained in younger patients when

number of oocytes vitrified was 24 [20]. Another study, evaluated the minimum number of mature oocytes to achieve at least one euploid blastocyst for transfer. The study found that the age of the woman was the most critical predictor for the likelihood of achieving one euploid blastocyst. Based on this model a patient of 37 years-old undergoing ART treatment using ejaculated sperm needs between 9 to 13 mature oocytes to obtain at least one euploid blastocyst to transfer [28]. Regarding the safety of the procedure, studies have analyzed the long term obstetric and perinatal outcomes associated with oocyte vitrification. An analysis of 165 pregnancies and 200 infants found that the mean birth weight and incidence of congenital abnormalities were similar in infants born following oocyte vitrification to those born from spontaneous conception or through standard ART treatment [17]. Another review of 936 infants, born following either slow-freezing or vitrification of oocytes, also reported a comparable incidence of congenital abnormalities [61]. A large study published in 2014 reported births of 1027 babies derived from vitrified-warmed oocytes and suggested that oocyte vitrification does not increase adverse obstetric and perinatal outcomes [19]. Thus, clinical outcomes using vitrified-warmed oocytes followed by IVF or ICSI appear to be similar to outcomes using fresh oocytes. However, these data were mainly reported for oocyte donation cycles and for standard ART cycles. Comparable data for women after cancer treatment who became pregnant and delivered a child after oocyte cryopreservation are not yet available.

Embryo cryopreservation in cancer patients

Oocytes obtained from OS can be fertilized using the partner's sperm, and cryopreserved for future use. The first pregnancy from cryopreserved embryos was reported in Australia in 1983 [93] and the first baby born after transfer of a cryopreserved-thawed blastocyst was announced in 1985 [21]. Initially, slow-freezing was the method used, but as with oocytes, this has now been replaced by vitrification. Embryo cryopreservation is the most established FP option for BC patients who have a male partner [39, 40] or for those women who are using donor sperm. Although this option is the most widely used globally, is not an option for couples who might have personal religious or moral objections. In addition, it is essential that the patient is informed and recognizes that any such embryos will require consent from both her and her partner for their subsequent use, and that may be problematical if the relationship is not continuing at the time of use [48]. Embryo

cryopreservation implies OS: as described above, recently studies have reported the use of OS protocols that can be started at anytime during the menstrual cycle [23]. Comparison of patients with and without cancer who underwent IVF and embryo cryopreservation have shown no difference in the number of collected oocytes, fertilization rates and number of live births, although patients with cancer had fewer good quality embryos [64]. Published studies have reported pregnancy outcomes comparable to those of non-oncological populations after IVF. Muñoz and collaborators performed a cohort study including 259 patients with early BC scheduled to receive chemotherapy (age 18 to 40 years old) divided into patients who wished to preserve their fertility (exposed group; n = 148), and underwent OS and chose to vitrify their oocytes, and patients with the same characteristics, but who did not want to preserve their fertility (non-exposed group; n = 111). The primary endpoint was disease free survival time and overall survival rate, with a follow-up of 5 years. Recurrences occurred in 9/148 women (6.1%) in the exposed group and 15/111 women (13.5%) in the non-exposed group, with no significant difference. The overall survival rates were comparable: 2/148 (1.4%) and 4/111 (3.6%) patients died, in exposed and non-exposed groups, respectively, therefore the authors concluded that ovarian stimulation in patients with early stage breast cancer appears safe in the long term [59]. A study published by Oktay and coauthors analysed OS with the concurrent use of letrozole in 131 women with BC with the purpose of FP via embryo freezing. Of the 131 women undergoing embryo cryopreservation, 33 come back to thaw their embryo and use in frozen embryo transfers. Post thaw survival rate of embryos was 98 (84.4%) and the mean number of embryos transferred was 1.97 ± 0.7 . They reported an overall clinical pregnancy per transfer of 65.0% (26 of 40), live birth per transfer of 45.0% (18 of 40), which is comparable to those in a non-cancer population undergoing ART treatment [66]. Table 1 displays published trials performed to assess ovarian performance in cancer, in which breast cancer disease was a predominant diagnosis.

Ovarian Tissue Cryopreservation (OTC)

OTC is a potential option for young women with breast cancer, though relatively infrequently used where oocyte vitrification is available. Although there are historic reports of ovarian transplantation in humans [62], the technique came to the fore following its successful development in the sheep,

266 where ovarian function and fertility were demonstrated after cryopreservation and
267 autotransplantation of ovarian cortical tissue [8, 32]. The first live birth was announced in 2004 [27],
268 and now more than 130 live births have been reported worldwide [30], demonstrating that this
269 strategy is viable in adults, although the success rate is unclear because the total number of attempts
270 performed is unknown. OTC involves the surgical removal (or dissection following oophorectomy
271 in many cases) and cryopreservation of the ovarian cortex. Later, upon completion of oncologic
272 treatment, the ovarian tissue can be thawed and transplanted back into the patient, either to
273 orthotopic (into the pelvic cavity; on the atrophic ovary) or heterotopic sites (outside of the pelvis;
274 subcutaneous regions such as the forearm) although only limited success has been reported from the
275 latter. It can be performed at any time during the menstrual cycle, there is no need for OS, and
276 therefore no delay in cancer treatment, and it results in storage of a large number of primordial
277 follicles, depending on the patient's age [27]. After reimplantation, ovarian function is expected to
278 be restored after 4-5 months, normally in more than 90% of patients. Regarding the freezing
279 procedure, slow freezing is most widely used: most centres use Gosden's protocol with
280 dimethylsulfoxide [60]. The efficiency of vitrification for freezing human ovarian tissue remains
281 controversial [1] but there have been two reports of births from vitrified and replaced ovarian tissue
282 [30]. Ovarian graft longevity is very variable but the woman's age is a crucial factor in determining
283 success, and many centres use an upper age limit of 35 years, in addition to criteria regarding risk of
284 infertility and chance of survival [3, 27]. Although, more than 130 live births have been reported
285 worldwide [30], there are still unresolved concerns, as substantial loss of primordial follicles is
286 known to occur after transplantation. This event seems to be related to the early hypoxia state that
287 characterizes the post-grafting period [52]. However, this loss of dormant follicles is accompanied
288 by an increase in the growing follicle population, suggesting a double mechanism of follicle death
289 and activation [53]. The greatest concern about this method is safety of the procedure relating to
290 that the replaced ovarian tissue might reimplant the cancer, therefore ovarian tissue should be
291 properly inspected, both by histology and immunohistochemistry (with additional molecular
292 analyses where possible) for malignant involvement of the ovarian tissue. This risk is however
293 considered low in early breast cancer [5].

Preimplantation genetic testing for monogenic disorder (PGT-M) to avoid BRCA transmission

The mean age at diagnosis of breast cancer for BRCA1 and BRCA2 mutation carriers is 43 and 47 years, respectively [96], but with a significant number of cases diagnosed before age 35. In BRCA1 carriers, the cancer incidence per year is 10/1000 in women between 20 and 29 years, 17/1000 between 30 and 39, and 20/1000 between 40 and 49 years. For BRCA2 carriers, the incidence peaks at age 40 to 49 (41/1000 cases per year) [54]. These women are therefore encouraged to undergo risk reducing salpingo-oophorectomy at ages 35-40 for BRCA1-carriers and between 40 and 45 for BRCA2-carriers [50]. PGT-M offers a valid option for BRCA-carriers women wishing to avoid transmission of the mutation to their offspring and being able to conceive an unaffected child. Preimplantation genetic testing in the human was successfully introduced in the late 1980s for fertile couples at risk of transmitting X chromosome-linked diseases to their children [35]. The process involves the aspiration of one or more cells from an embryo generated through IVF, subsequent genetic analysis, and the transfer into the uterus of only unaffected embryos [11, 35]. As stated earlier, the evolution of pre-implantation genetic assessment started with the analysis of limited number of chromosomes using the fluorescence in situ hybridization (FISH) technology in the late 1980s [11, 35]. It was soon replaced by analysis of the whole chromosome set by using different genetic platforms, such as metaphase Comparative Genomic Hybridization (CGH), array based Comparative Genomic Hybridization (aCGH), single nucleotide polymorphism (SNP) microarray, and quantitative polymerase chain reaction (qPCR). At present, the most advanced technique is Next Generation Sequencing (NGS), which refers to a DNA sequencing technology that enables sequencing of millions of small DNA fragments in unison. NGS has revolutionized genomic research studies, and is currently the gold standard for the analysis of monogenic diseases or single single gene mutations [84]. As an autosomal dominant, women with a BRCA mutation have a 50% chance of transferring it to their offspring. BRCA1 and BRCA2 are members of the ATM (ataxia telangiectasia mutation) protein family, involved in DNA double strand damage detections and repairs. Loss of ATM function in human and mouse causes defects in DNA repair and cell cycle checkpoint control and thus predisposes to cancers. BRCA1 is also highly expressed in germ cells and blastocysts, suggesting a possible role in gametogenesis and embryogenesis. In

the oocytes of primordial follicles in BRCA mutation carriers, it has been suggested that DNA damage may accumulate over time: this may lead to loss of some follicles, with a reduction in the ovarian reserve. This correlation has been demonstrated in mice model, where BRCA1 mutation is associated with lower primordial follicle counts and AMH levels compared to normal controls [92] and there are data suggesting the same in women, for BRCA1 but not BRCA2 [65, 71, 92]. Women with BRCA mutations may show a reduced ovarian response to OS [46] although not all studies have confirmed this [85]. With the PGT-M technique, embryos cultured in vitro are genetically tested for the presence of the mutation, in order to transfer only BRCA negative embryos to the uterus. Couples undergoing PGT-M are usually fertile but they have to undergo IVF treatment, which can be costly and stressful. These couples also have to face the possibility that all embryos might be affected, and that the transfer of an unaffected embryo may not lead to a successful pregnancy. In 2003, despite uncertainties about prospective improvements and therapeutic opportunity, the European Society of Human Reproduction and Embryology (ESHRE) ethics taskforce considered genetic testing acceptable for hereditary conditions and multifactorial diseases such as BC or other cancer dispositions [86]. A major benefit compared to the alternative approach of prenatal testing is the avoidance of consideration of termination of an otherwise viable pregnancy. It is important to recognize that PGT-M is not a therapy, but only a selection tool. As an autosomal dominant condition, half of the embryos will be expected to test positive for the relevant BRCA mutation and thus will be discarded. As the number of available embryos will decline with the woman's age and the number of oocytes collected, it seems more appropriate only in young BC patients. As discussed above, being a carrier of a BRCA mutation may also reduce the number of embryos available for testing. Moreover, for PGT-M a physically demanding in vitro fertilization treatment is required regardless of couple's fertility, and OS is necessary, which can delay cancer treatment [39, 40]. Opinion studies among women affected by BC have shown that the majority, after being informed about PGT-M, are in favour of offering PGT-M for BRCA1 and BRCA2 mutations, although only a minority would consider this option for themselves [24, 67]. PGT-M for BRCA mutations is growing; a survey of 1081 BRCA mutation carriers highlighted that patients are keen to have reproductive counseling, with more than 50% stating that PGT-M should be offered.

The most frequently quoted reason in considering PGT-M was, in all categories of couples, to protect their future child from the physical and psychological impact of the BRCA mutation [15].

Conclusion

FP is a rapidly developing area of medicine, and the provision of information to patients facing the loss of fertility through treatment for cancer and other conditions has become standard of care. Women should be informed not only about advantages of oocyte and embryo cryopreservation, as established technology that will contribute to achieve a pregnancy after cancer, but also about the general risks, cost and effectiveness of the procedures to reach a shared decision. Reproductive decision-making regarding PGT-M is complex for BRCA mutation carriers. For some couples, the emotional impact of the decision is substantial and long-lasting, therefore reproductive and dynamic counselling over time is crucial, considering that a woman's aspirations may change with age. All women about to receive chemotherapy for a newly diagnosed BC should receive proper and complete oncofertility counselling regarding the possible gonadotoxic risk and potential approaches for FP, to allow them to take fully informed decisions about the proposed therapy and its long-term consequences. This requires as a minimum the development of optimized communication between specialities, with referral to reproductive medicine clinics for ART becoming an integrated part of cancer care. The development of national and international registries is required to monitor the techniques used, the success rates achieved and the long-term follow-up of children born from these procedures.

Conflict of interest. RS declares that he has no conflict of interest. RAA is past coordinator of the ESHRE Special Interest Group in Fertility Preservation.

Compliance with ethical standards

Human and animal rights. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and with the 1964 Helsinki declaration and its later amendments. For this type of study, formal consent is not required.

Funding and role of the funding body RAA is at the MRC Centre for Reproductive Health, which is funded by MRC Centre grant MR/N022556/1

381

382 **References**

383

- 384 [1] R. Abir, B. Fisch, N. Fisher, N. Samara, G. Lerer-Serfaty, R. Magen, M. Herman-Edelstein, A.
 385 Ben-Haroush, A. Stein, R. Orvieto, Attempts to improve human ovarian transplantation outcomes
 386 of needleimmersed vitrification and slow-freezing by host and graft treatments. *Journal of Assisted*
 387 *Reproduction and Genetics* 2017, 34; 633-644.
- 388 [2] M.E. Abusief, S.A. Missmer, E.S. Ginsburg, J.C. Weeks, A.H. Partridge, The effects of
 389 paclitaxel, dose density, and trastuzumab on treatment-related amenorrhea in premenopausal
 390 women with breast cancer. *Cancer* (2010) 116:791-8.
- 391 [3] R.A. Anderson, R. T. Mitchell, T. W. Kelsey, N. Spears, E. E. Telfer, W. H. Wallace, Cancer
 392 treatment and gonadal function: experimental and established strategies for fertility preservation in
 393 children and young adults. *Lancet Diabetes Endocrinol* 2015; 3:556-567
- 394 [4] R.A. Anderson, D.H. Brewster, R. Wood, S. Nowell, C. Fischbacher, T.W. Kelsey and W.H.B.
 395 Wallace, The impact of cancer on subsequent chance of pregnancy: a population-based analysis.
 396 *Hum Reprod* 2018; 33: 1281-1290.
- 397 [5] R.A. Anderson, D. T. Baird, The development of ovarian tissue cryopreservation in Edinburgh:
 398 Translation from a rodent model through validation in a large mammal and then into clinical
 399 practice. *Acta Obstet Gynecol Scand* 2019; 98: 545-549.
- 400 [6] R.A. Anderson, D.A. Cameron, Pretreatment serum anti-mullerian hormone predicts long-term
 401 ovarian function and bone mass after chemotherapy for early breast cancer. *J Clin Endocrinol*
 402 *Metab* 2011; 96: 1336-1343;
- 403 [7] R.A. Anderson, M. Rosendahl, T.W. Kelsey, D.A. Cameron, Pretreatment anti-Mullerian
 404 hormone predicts for loss of ovarian function after chemotherapy for early breast cancer. *Eur J*
 405 *Cancer* 2013; 49: 3404-3411;
- 406 [8] D.T. Baird, R. Webb, B. K. Campbell, L. M. Harkness, R. G Gosden, Long-term ovarian
 407 function in sheep after ovariectomy and transplantation of autografts stored at -196 C.
 408 *Endocrinology* 1999; 140: 462-471.
- 409 [9] H. Bar-Joseph, I. Ben-Aharon, S. Rizel, S.M. Stemmer, M. Tzabari, R. Shalgi, Doxorubicin-
 410 induced apoptosis in germinal vesicle (GV) oocytes. *Reprod Toxicol* (2010) 30:566-72.
 411
- 412 [10] A. Bielanski, S. Nadin-Davis, T. Sapp, C. Lutze-Wallace, Viral contamination of embryos
 413 cryopreserved in liquid nitrogen. *Cryobiology* 40 (2000) 40:110-116.
 414
- 415 [11] P. Braude, S. J. Pickering, F. Flinter, C. M. Ogilvie, Preimplantation genetic diagnosis. *Nat*
 416 *Rev Genet* 2002 Dec ;3 (12) :941-953.
 417
- 418 [12] F. Bray, J. Ferlay, I. Soerjomataram, R.L. Siegel, L.A. Torre LA, A. Jemal, Global cancer
 419 statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185
 420 countries. *CA Cancer J Clin* 2018, 68 (6): 394-424.

- 421 [13] D. Brison, R. Cutting, H. Clarke, M. Wood, ACE consensus meeting report: oocyte and
422 embryo cryopreservation Sheffield 17.05.11. Hum Fertil (2012); 15: 69-74.
- 423 [14] E.R. Cardozo, A.P. Thomson, A.E. Karmon, K.A. Dickinson, D.L. Wright, M.E. Sabatini,
424 Ovarian stimulation and in-vitro fertilization outcomes of cancer patients undergoing fertility
425 preservation compared to age matched controls: a 17-year experience. J Assist Reprod Genet. 2015;
426 32 (4): 587-96.
- 427 [15] J.L. Chan, L. N. C. Johnson, M. D. Sammel, L. Giovanni, C. Voong, S. M. Domchek, C. R.
428 Gracia, Reproductive decision-making in women with BRCA 1-2 mutations. J Genet Couns 2017;
429 26: 594-603.
- 430 [16] C. Chen, Pregnancy after human oocyte cryopreservation. Lancet (1986) 1:884-886.
- 431 [17] R.C. Chian, J. Y. Huang, S. L. Tan, E. Lucena, A. Saa, A. Rojas, L. A. Ruvalcaba Castellon,
432 M. I. Garcia Amador, J. E. Montoya Sarmiento, Obstetric and perinatal outcome in 200 infants
433 conceived from vitrified oocytes. Reprod Biomed Online (2008); 16: 608-610.
- 434 [18] A. Cobo, C. Diaz, Clinical application of oocyte vitrification: a systematic review and meta-
435 analysis of randomized controlled trials. Fertil Steril (2011); 96: 277-285.
- 436 [19] A. Cobo, V. Serra, N. Garrido, I. Olmo, A. Pellicer, J. Remohí, Obstetric and perinatal
437 outcome of babies born from vitrified oocytes. Fertil Steril. 2014 Oct;102(4):1006-1015.
- 438 [20] A. Cobo, J. García-Velasco, J. Domingo, A. Pellicer, J. Remohí, Elective and Onco-fertility
439 preservation: factors related to IVF outcomes. Hum Reprod 2018 Dec 1;33(12): 2222-2231.
- 440 [21] J. Cohen, R. F. Simons, C. B. Fehillity, S. B. Fishel, R. G. Edwards, J. Hewitt, G. F. Rowland,
441 P. C. Steptoe, J. M. Webster, Birth after replacement of hatching blastocyst cryopreserved at
442 expanded blastocyst stage. Lancet (1985)16: 647.
- 443 [22] S.S Coughlin, Epidemiology of Breast Cancer in Women. Adv Exp Med Biol. 2019;1152:9-29.
- 444 [23] R.B. Danis, N. Pereira, R. T. Elias, Random start ovarian stimulation for oocyte or embryo
445 cryopreservation in women desiring fertility preservation prior to gonadotoxic cancer therapy. Curr
446 Pharm Biotechnol 2017, 18 (8): 609-13.
- 447 [24] C. Dekeuwer, S. Bateman, Much more than a gene: hereditary breast and ovarian cancer,
448 reproductive choices and family life. Med Health Care Philos 2013; 16: 231-244.
449
- 450 [25] I. Demeestere, F. Moffa, F. Peccatori, C. Poirot, E. Shalom-Paz, Multiple approaches for
451 individualized fertility protective therapy in cancer patients. Obstet Gynecol Int. 2012; 2012:
452 961232.
- 453 [26] J. Domingo, V. Guillén, Y. Ayllón, M. Martínez, E. Muñoz, A. Pellicer, J.A. Garcia-Velasco.,
454 Ovarian response to controlled ovarian hyperstimulation in cancer patients is diminished even
455 before oncological treatment. Fertil Steril. 2012;97(4):930-4.

- 456 [27] J. Donnez, M. M. Dolmans, D. Demylle, P. Jadoul, C. Pirard, J. Squifflet, B. Martinez-Madrid,
457 A. van Langendonckt, Livebirth after orthotopic transplantation of cryopreserved ovarian tissue.
458 *Lancet* 2004, 364 1405-1410.
- 459 [28] S.C. Esteves, J.F. Carvalho, F.C. Bento, J.Santos, A Novel Predictive Model to Estimate the
460 Number of Mature Oocytes Required for Obtaining at Least One Euploid Blastocyst for Transfer in
461 Couples Undergoing in vitro Fertilization/Intracytoplasmic Sperm Injection: The ART Calculator.
462 *Front Endocrinol (Lausanne)* 2019 Feb 28; 10:99.
- 463 [29] B.J. Fuller, J.E. Hunter, A.G. Bernard, J.J. McGrath, P. Curtis, A. Jackson, The permeability of
464 unfertilised oocytes to 1,2-propanediol: a comparison of mouse and human cells. *CryoLetters*
465 (1992) 13: 287-292.
- 466 [30] S.E. Gellert, S.E. Pors, S.G. Kristensen, A.M. Bay-Bjorn, E. Ernst, C. Yding Andersen,
467 Transplantation of frozen-thawed ovarian tissue: an update on worldwide activity published in peer-
468 reviewed papers and on the Danish cohort. *J Assist Reprod Genet* 2018; 35: 561-570.
- 469 [31] R.B. Gilchrist, M. Lane M, J.G. Thompson, Oocyte-secreted factors: regulators of cumulus cell
470 function and oocyte quality. *Hum Reprod Update* 2008; 14: 159-177.
- 471 [32] R.G. Gosden, D.T. Baird, J.C. Wade, R. Webb, Restoration of fertility to oophorectomized
472 sheep by ovarian autografts stored at -196°C. *Human Reproduction* 1994; 9: 597-603.
- 473 [33] R. Gualtieri, M. Iaccarino, V. Mollo, M. Prisco, S. Iaccarino, R. Talevi, Slow cooling of
474 human oocytes: ultrastructural injuries and apoptotic status. *Fertil Steril* 2009; 91(4): 1023- 1034.
- 475 [34] T. Haahr T, S.C. Esteves, P. Humaidan, Individualized controlled ovarian stimulation in
476 expected poor-responders: an update. *Reprod Biol Endocrinol* 2018, 16:20.
- 477
478 [35] A.H. Handyside, E. H. Kontogianni, K. Hardy K, R. M. Winston, Pregnancies from biopsied
479 human preimplantation embryos sexed by Y-specific DNA amplification. *Nature*. 1990;344:768-
480 770.
- 481
482 [36] A.J. Hsueh, K. Kawamura, Y. Cheng, B. C. Fauser, Intraovarian control of early
483 folliculogenesis. *Endocr Rev* 2015; 36:1-24.
- 484
485 [37] L.N. Johnson, K.E. Dillon KE, M.D. Sammel, B.L. Efymowm, M.A. Mainigi, A. Dokras, C.R.
486 Gracia, Response to ovarian stimulation in patients facing gonadotoxic therapy. *Reprod Biomed*
487 *Online*. 2013;26(4):337-44.
- 488
489 [38] A. Jones, J. Van Blerkom, P. Davis, A. A. Toledo, Cryopreservation of metaphase II human
490 oocytes effects mitochondrial membrane potential: implications for developmental competence.
491 *Hum Reprod* (2004) ;19 (8):1861- 1866.
- 492 [39] S.S. Kim, J. Klemp, C. Fabian, Breast cancer and fertility preservation. *Fertil Steril* (2011a)
493 95:1535-1543.
- 494 [40] S.S. Kim, K. Oktay, C. Gracia, S. Lee, C. Morse, J.E. Mersereau, Which patients pursue
495 fertility preservation treatments? A multicenter analysis of the predictors of fertility preservation in
496 women with breast cancer. *Fertil Steril* (2012) 97:671-676.

- [41] S.Y. Kim, S.K. Kim, J.R. Lee, T.K. Woodruff, Toward precision medicine for preserving fertility in cancer patients: existing and emerging fertility preservation options for women. *J Gynecol Oncol* (2016) 27(2): e22.
- [42] J.M. Knopman, N. Noyes, S. Talebian, L.C. Krey, J.A. Grifo, F. Licciardi, Women with cancer undergoing ART for fertility preservation: a cohort study of their response to exogenous gonadotropins. *Fertil Steril*. 2009;91(4 Suppl):1476–8.
- [43] L. Kuleshova, L. Gianaroli, C. Magli, A. Ferraretti, A. Trounson, Birth following vitrification of a small number of human oocytes: case report. *Human Reproduction* (1999) 14: 3077-3079.
- [44] M. Lambertini, N. Kroman, L. Ameye, O. Cordoba, A. Pinto, G. Benedetti, M.B. Jensen, S. Gelber, M. Del Grande, M. Ignatiadis, E. de Azambuja, M. Paesmans, F.A. Peccatori and H.A. Jr. Azim, Long-term Safety of Pregnancy Following Breast Cancer According to Estrogen Receptor Status. *J Natl Cancer Inst* 2018; 110: 426-429.
- [45] M. Lambertini, H.C.F. Moore, R.C.F. Leonard, S. Loibl, P. Munster, M. Bruzzzone, L. Boni, J.M. Unger, R.A. Anderson, K. Mehta, S. Minton, F. Poggio, K.S. Albain, D.J.A. Adamson, B. Gerber, A. Cripps, G. Bertelli, S. Seiler, M. Ceppi, A.H. Partridge and L. Del Mastro, Gonadotropin-Releasing Hormone Agonists During Chemotherapy for Preservation of Ovarian Function and Fertility in Premenopausal Patients With Early Breast Cancer: A Systematic Review and Meta-Analysis of Individual Patient-Level Data. *J Clin Oncol* 2018a; 36: 1981-1990.
- [46] M. Lambertini, O. Goldrat, A. R. Ferreira, J. Dechene, H. A. Jr. Azim, J. Desir, A. Delbaere, M. D. t'Kint de Roodenbeke, E. de Azambuja, M. Ignatiadis, I. Demeestere, Reproductive potential and performance of fertility preservation strategies in BRCA-mutated breast cancer patients. *Ann Oncol*. 2018 Jan 1;29(1):237-243.
- [47] J.A. Lee, L. Sekhon, L. Grunfeld, A.B. Copperman, In-vitro maturation of germinal vesicle and metaphase I eggs prior to cryopreservation optimizes reproductive potential in patients undergoing fertility preservation. *Curr Opin Obstet Gynecol*. 2014;26(3):168-73.
- [48] S.J. Lee, L.R. Schover, A.H. Partridge, P. Patrizio, W.H. Wallace, K. Hagerty, L.N. Beck, L.V. Brennan, K. Oktay, American Society of Clinical Oncology recommendations on fertility preservation in cancer patients. *J Clin Oncol* (2006); 24: 2917-2931.
- [49] J.M. Letourneau, E.E Ebbel, P.P. Katz, K.H Oktay, C.E. McCulloch, W.Z Ai, A.J. Chien, M.E. Melisko, M.I. Cedars, M.P. Rosen, Acute ovarian failure underestimates age-specific reproductive impairment for young women undergoing chemotherapy for cancer. *Cancer* 2012a; 118:1933-1939.
- [50] E.K. Lewis, K.H. Lu, A.M. Klimczak, S.C. Mok, Recommendations and Choices for BRCA Mutation Carriers at Risk for Ovarian Cancer: A Complicated Decision. *Cancers (Basel)*. 2018 Feb; 10(2): 57.
- [51] A.W. Loren, P. B. Mangu, L. N. Beck, L. Brennan, A. J. Magdalinski, A. H. Partridge, G. Quinn, W. H. Wallace, K. Oktay, Fertility preservation for patients with cancer: American Society of Clinical Oncology clinical practice guideline update. *J Clin Oncol* 2013; 31:2500-2510.
- [52] D.D. Manavella, L. Cacciottola, S. Pomme, C.M. Desmet, B.F. Jordan, J. Donnez, C.A. Amorim, M.M. Dolmans, Two-step transplantation with adipose tissue-derived stem cells increases

follicle survival by enhancing vascularization in xenografted frozen-thawed human ovarian tissue. Hum Reprod. 2018;33:1107-16).

[53] R. Masciangelo, C. Hossay, J. Donnez, M.M Dolmans, Does the Akt pathway play a role in follicle activation after grafting of human ovarian tissue? Reprod BioMed Online 2019; 39(2):196-8.

[54] N. Mavaddat, S. Peock, D. Frost, S. Ellis, R. Platte, E. Fineberg, D. G. Evans, L. Izatt, R.A. Eeles, J. Adlard et al, Cancer risks for BRCA1 and BRCA2 mutation carriers: results from prospective analysis of EMBRACE. J Natls Cancer Inst 2013 Jun 5; 105 (11): 812-822.

[55] D. Meirrow, H. Biederman, R.A. Anderson, W.H. Wallace, Toxicity of chemotherapy and radiation on female reproduction. Clin Obstet Gynecol (2010) 53:727-39.

[56] D.F. Merlo, M. Ceppi, R. Filiberti, V. Bocchini, A. Znaor, M. Gamulin, M. Primic-Žakelj, P. Bruzzi, C. Bouchardy, A. Fucic, W.G. Airtum, Breast cancer incidence trends in European women aged 20-39 years at diagnosis. Breast Cancer Res Treat. 2012 Jul;134(1):363-70.

[57] N. Michaan, G. Ben-David, D. Ben-Yosef, B. Almog, A. Many, D. Pauzner, J.B. Lessing, A. Amit, F. Azem, Ovarian stimulation and emergency in vitro fertilization for fertility preservation in cancer patients. Eur J Obstet Gynecol Reprod Biol. 2010;149 (2):175-7.

[58] S. Morgan, R.A. Anderson, C. Gourley, W.H. Wallace and N. Spears, How do chemotherapeutic agents damage the ovary? Hum Reprod Update 2012; 18: 525-535.

[59] E. Muñoz, J. Domingo, G. De Castro, I. Lorenzo, J. A García-Velasco, J. Bellver, A. Pellicer, N. Garrido, Ovarian stimulation for oocyte vitrification does not modify disease-free survival and overall survival rates in patients with early breast cancer. Reprod Biomed Online 2019 Jul 10. pii: S1472-6483(19)30633-9

[60] H. Newton, J. Fisher, J. R. Arnold, D. E. Pegg, M. J. Faddy, R. G. Gosden, Permeation of human ovarian tissue with cryoprotective agents in preparation for cryopreservation. Human Reproduction 1998, 13; 376-380.

[61] N. Noyes, E. Porcu, A. Borini, Over 900 cryopreservation babies born with no apparent increase in congenital anomalies. Reprod Biomed Online (2009); 18: 769-776.

[62] D. Nugent, D. Meirrow, P. F. Brook, Y. Aubard, R. G. Gosden, Transplantation in reproductive medicine: previous experience, present knowledge and future prospects. Hum Reprod Update 1997; 3: 267-280.

[63] ONS. Cancer survival in England: one-year and five-year survival for 21 common cancers, by sex and age. Office for National Statistics, 2010.

[64] K. Oktay, A. P. Cil, H. Bang, Efficiency of oocyte cryopreservation: a meta-analysis. Fertil Steril 2006 Jul; 86 (1):70-80.

[65] K. Oktay, J. Y. Kim, D. Barad, S. N. Babayev, Association of BRCA1 mutations with occult primary ovarian insufficiency: a possible explanation for the link between infertility and breast/ovarian cancer risks. J Clin Oncol 2010; 28:240244.

585

586 [66] K. Oktay, V. Turan, G. Bedoschi, F.S. Pacheco, F. Moy, Fertility preservation success
 587 subsequent to concurrent aromatase inhibitor treatment and ovarian stimulation in women with
 588 breast cancer. *J Clin Oncol.* 2015; 33(22):2424-9.

589 [67] E. Ormondroyd, L. Donnelly, C. Moynihan, C. Savona, E. Bancroft, D. G. Evans, R. Eeles, S.
 590 Lavery, M. Watson, Attitudes to reproductive genetic testing in women who had a positive BRCA
 591 test before having children: a qualitative analysis. *Eur J Hum Genet* 2012; 20:4-10.
 592

593 [68] S.J. Paynter, A. Cooper, L. Gregory L, B. J. Fuller, R.W. Shaw, Permeability characteristics of
 594 human oocytes in the presence of the cryoprotectant dimethylsulphoxide. *Hum Reprod* (1999) 14:
 595 2338-2342.

596 [69] M. Peate, B. Meiser, B.C. Cheah, C.Saunders, P. Butow, B. Thewes, R. Hart, K.A. Phillips, M.
 597 Hickey, M. Friedlander, Making hard choices easier: a prospective, multicentre study to assess the
 598 efficacy of a fertility-related decision aid in young women with early-stage breast cancer. *Br*
 599 *J Cancer.* 2012 Mar 13;106(6):1053-61.

600 [70] J.A. Petrek, M.J. Naughton, L.D. Case, E.D. Paskett, E.Z. Naftalis, S.E. Singletary, P.
 601 Sukumvanich, Incidence, time course, and determinants of menstrual bleeding after breast cancer
 602 treatment. A prospective study. *J Clin Oncol* 2006; 24:1045-1051.
 603

604 [71] K. A. Phillips, I. M. Collins, R. L. Milne, S. A. McLachlan, M. Friedlander, M. Hickey, C.
 605 Stern, J. L. Hopper, R. Fisher, G. Kannemeyer, S. Picken, C. D. Smith, T. W. Kelsey, R. A.
 606 Anderson, K. Cuningham, Consortium for Research into Familial Breast Cancer Anti-Mullerian
 607 hormone serum concentrations of women with germline BRCA1 or BRCA2 mutations. *Hum*
 608 *Reprod* 2016; 31: 1126-1132.

609 [72] S.J. Pickering, P.R. Braude, M. H. Johnson, A. Cant A, J. Currie, Transient cooling to room-
 610 temperature can cause irreversible disruption of the meiotic spindle in the human oocyte. *Fertil*
 611 *Steril* (1990) 54:102-108.

612 [73] E. Porcu, R. Fabbri, R. Seracchioli, P. M. Ciotti, O. Magrini, C. Flamigni, Birth of a healthy
 613 female after intracytoplasmic sperm injection of cryopreserved human oocytes. *Fertil Steril* (1997)
 614 68:724-726.

615 [74] Practice Committees of American Society for Reproductive Medicine; Society for Assisted
 616 Reproductive Technology. Mature oocyte cryopreservation: a guideline. *Fertil Steril* (2013) Jan;
 617 99(1): 37-43. Epub 2012 Oct 22.

618 [75] M.M. Quinn, H. Cakmak, J. M. Letourneau, M. I. Cedars, M. P. Rosen, Response to ovarian
 619 stimulation is not impacted by a breast cancer diagnosis. *Hum Reprod.* 2017 Mar 1;32(3):568-574.

620 [76] R.B. Quintero, A. Helmer, J.Q. Huang, L.M. Westphal, Ovarian stimulation for fertility
 621 preservation in patients with cancer. *Fertil Steril.* 2010;93(3):865-8.

- [77] L. Rienzi, S. Romano, L. Albricci, R. Maggiulli, A. Capalbo, E. Baroni, S. Colamaria, F. Sapienza, F. M. Ubaldi, Embryo development of fresh 'versus' vitrified metaphase II oocytes after ICSI: a prospective randomized sibling-oocyte study. *HumReprod* (2010);2 5: 66-73.
- [78] L. Rienzi, C. Gracia, R. Maggiulli, A. R. LaBarbera, D. J. Kaser, F. M. Ubaldi, S. Vanderpoel, C. Racowsky, Oocyte, embryo and blastocyst cryopreservation in ART: systematic review and meta-analysis comparing slow-freezing versus vitrification to produce evidence for the development of global guidance. *Hum Reprod Update* (2017). 23:139-155.
- [79] A.D. Robertson, S.A. Missmer, E.S. Ginsburg, Embryo yield after in vitro fertilization in women undergoing embryo banking for fertility preservation before chemotherapy. *Fertil Steril*. 2011;95(2): 588-91.
- [80] K.J. Ruddy, S.I. Gelber, R.M. Tamimi, E.S. Ginsburg, L. Schapira, S.E. Come, F.V. Borges, M.E. Meyer and A.H. Partridge, Prospective study of fertility concerns and preservation strategies in young women with breast cancer. *J Clin Oncol*. 2014;32:1151-1156.
- [81] L.R. Schover, M. van der Kaaij, E. van Dorst, C. Creutzberg, E. Huyghe, C.E. Kiserud, Sexual dysfunction and infertility as late effects of cancer treatment. *EJC Suppl EJC Off J EORTC Eur Organ Res Treat Cancer Al*. (2014) 12(1):41-53.
- [82] R. Sciorio, K.J. Thong, S.J. Pickering, Single blastocyst transfer (SET) and pregnancy outcome of day 5 and day 6 human blastocysts vitrified using a closed device *Cryobiology*. 2018 Oct; 84: 40-45. Epub 2018 Aug 10.
- [83] R. Sciorio, K. J. Thong, S. J. Pickering, Increased pregnancy outcome after day 5 versus day 6 transfers of human vitrified-warmed blastocysts. *Zygote*. 2019 Oct;27(5):279-284.
- [84] K. Sermon, Novel technologies emerging for preimplantation genetic diagnosis and preimplantation genetic testing for aneuploidy. *Expert Rev Mol Diagn*. 2017;17:71-82.
- [85] M. Shapira, H. Raanani, D. Meirow, IVF for fertility preservation in breast cancer patients- efficacy and safety issues. *J Assist Reprod Genet* 2015a; 32:1171-1178.
- [86] F. Shenfield, G. Pennings, P. Devroey, C. Sureau, B. Tarlatzis, J. Cohen, ESHRE ethics task force. Taskforce 5: preimplantation genetic diagnosis. *Hum Reprod* 2003;18: 649-651.
- [87] N. Spears, F. Lopes, A. Stefansdottir, V. Rossi, M. De Felici, R.A Anderson and F.G Klinger, Ovarian damage from chemotherapy and current approaches to its protection. *Hum Reprod Update* 2019 Oct 10. pii: dmz027. doi: 10.1093/humupd/dmz027.
- [88] V. Stearns, B. Schneider, N.L. Henry NL, D.F. Hayes, D. A Flockhart, Breast cancer treatment and ovarian failure: risk factors and emerging genetic determinants. *Nat Rev Cancer* 2006;6: 886-893.
- [89] H.C. Su, C. Haunschild, K. Chung, S. Komrokian, S. Boles, M.D. Sammel, A. DeMichele, Prechemotherapy antimullerian hormone, age, and body size predict timing of return of ovarian function in young breast cancer patients. *Cancer* 2014; 120: 3691-3698.

- 664 [90] H.I. Su, K. Chung, M.D. Sammel, C.R. Gracia, A. DeMichele, Antral follicle count provides
 665 additive information to hormone measures for determining ovarian function in breast cancer
 666 survivors. *Fertil Steril* 2011; 95: 1857-1859.
- 667 [91] W. Tarumi, N. Suzuki, N. Takahashi, Y. Kobayashi, K. Kiguchi, K. Sato, B. Ishizuka, Ovarian
 668 toxicity of paclitaxel and effect on fertility in the rat. *J Obstet Gynaecol Res* (2009) 35:414-20.
- 669 [92] Titus S, Li F, Stobezki R, Akula K, Unsal E, Jeong K, Dickler M, Robson M, Moy F,
 670 Goswami S, Oktay K. Impairment of BRCA1-related DNA doublestrand break repair leads to
 671 ovarian aging in mice and humans. *Sci Transl Med* 2013 Feb 13; 5 (172): 172ra21.
- 672 [93] A. Trounson, L. Mohr, Human pregnancy following cryopreservation, thawing and transfer of
 673 an eight-cell embryo. *Nature* (1983); 305: 707-709.
- 674 [94] V.Turan, M. M. Quinn, N. Dayioglu, M. P. Rosen, K. Oktay, The impact of malignancy on
 675 response to ovarian stimulation for fertility preservation: a meta-analysis. *Fertil*
 676 *Steril*. 2018 Dec;110(7):1347-1355.
- 678 [95] D. M. van der Kolk, G. H. de Bock, B. K. Leegte, M. Schaapveld, M. J. Mourits, J. de Vries,
 679 A. H. van der Hout, J. C. Oosterwijk, Penetrance of breast cancer, ovarian cancer and contralateral
 680 breast cancer in BRCA1 and BRCA2 families: high cancer incidence at older age. *Breast Cancer*
 681 *Res Treat* 2010; 124: 643-651.
- 682 [96] P. Vanderzwalmen, N.H. Zech, Y. Prapas, Y. Panagiotidis, A. Papatheodorou, B. Lejeune, D.
 683 Jarenˆo, S. Vanderzwalmen, F. Ectors, Closed carrier device: a reality to vitrify oocytes and
 684 embryos in aseptic conditions. *Gynecol Obstet Fertil* (2010) 38: 541-546.
- 685 [97] C. Vincent, S. J. Pickering, M. H. Johnson, The hardening effect of dimethylsulphoxide on the
 686 mouse zona pellucida requires the presence of an oocyte and is associated with a reduction in the
 687 number of cortical granules present. *J Reprod Fertil* (1990) 89:253-259.
- 689 [98] J. Wise, UK lifts ban on frozen eggs. *BMJ* (2000) Feb 5; 320 (7231):334.
- 690 [99] L.Y. Yan, J. Yan, J. Qiao, P.L. Zhao, P. Liu, Effects of oocyte vitrification on histone
 691 modifications. *Reprod Fertil Dev* (2010) ;22 (6): 920-925.

Fertility preservation and preimplantation genetic assessment for women with breast cancer

Romualdo Sciorio^{1*}, Richard A. Anderson²

- 1. Edinburgh Assisted Conception Programme, EFREC, Royal Infirmary of Edinburgh, 51 Little France Crescent, Edinburgh, UK.**
- 2. MRC Centre for Reproductive Health, The Queen's Medical Research Institute, Edinburgh BioQuarter, The University of Edinburgh, 47 Little France Crescent, Edinburgh, UK.**

***Corresponding author. E-mail address: sciorioromualdo@hotmail.com (R. Sciorio).**

Figure 1: Potential targets of chemotherapeutic damage within the ovary. **(A)** Chemotherapeutic agents could be directly affecting the resting pool of primordial follicles or the growing follicle population. As growing follicles inhibit the recruitment of primordial follicles, the loss of this growing population will lead to increased activation of primordial follicles and so loss of that reserve. **(B)** Chemotherapeutic agents could be directly targeting the oocyte or the somatic cells. Oocyte death would result from death of the follicular somatic cells, as the oocyte is dependant on these for its survival.

Reprinted with permission from Morgan et al, 2012 [45].




Figure 2. (A) Probability of pregnancy after cancer diagnosis in women with breast cancer (red) compared to matched population controls (blue). Table under the panel indicate the number of women at each 10 year interval. (B) Hazard ratio for first pregnancy after breast cancer diagnosis by period of diagnosis.

Reprinted from Anderson et al, 2018 [4], with permission.




Table 1: Compares IVF outcomes in cancer patients versus non cancer patients

Studies	Number of Patients ¹	% Breast Cancer ²	Mature Oocytes ³		Fertilized 2 PN	
			Cancer	Control	Cancer	Control
Cardozo et al. [14]	63	41 (65%)	12.4	10.9	6.6	7.1
Domingo et al. [26]	208	142 (69%)	10.5	12.4	N/A	N/A
Knopman et al. [42]	26	10 (38%)	14	12	N/A	N/A
Michaan et al. [57]	22	12 (55%)	8.8	8.8	5.4	5
Robertson et al. [79]	38	16 (42%)	12	14	6	7
Quintero et al. [76]	50	28 (56%)	11.5	13	6.8	7.4
Johnson et al. [37]	50	29 (58%)	12.4	11.7	5.4	6

1 Number of cancer patients included in trial

2 Number and percentage of breast cancer patients included in study

3 Mean number of oocytes collected for cancer patients and control patients.